

Professional Nursing Service

Critical Care/Intensive Care Checklist

Name: _____

Date: _____

Years of Experience: _____

Directions for completing skills checklist:

The following is a list of equipment and/or procedures performed in rendering care to patients. Please indicate your level of experience/proficiency with each area and, where applicable, the types of equipment and/or systems you are familiar with. Use the following key as a guideline:

- A) Theory Only/No Experience--Didactic instruction only, no hands on experience
- B) Limited Experience--Knows procedure/has used equipment, but has done so infrequently or not within the last six months
- C) Moderate Experience--Able to demonstrate equipment/procedure, performs the task/skill independently with only resource assistance needed.
- D) Proficient/Competent--Able to demonstrate/perform the task/skill proficiently without any assistance and can instruct/teach.

A. CARDIOVASCULAR

	A	B	C	D
1. Assessment				
a. Abnormal heart sounds/murmurs/arrhythmias	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Auscultation (rate, rhythm)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Blood pressure/non-invasive	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Doppler	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Pulses/circulation checks	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2. Interpretation of lab results				
a. Cardiac enzymes & isoenzymes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Coagulation studies	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3. Equipment & procedures				
a. Assist with				
(1) Arterial line insertion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Central line insertion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Open chest emergency	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) PA catheter/Swan-Ganz insertion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) Pericardiocentesis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(6) Transesophageal echocardiogram	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Automatic internal cardioverter defibrillator	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Cardioversion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. CAVH-D	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Hemodynamic monitoring				
(1) Cardiac index	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Cardiac output	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) CVP monitoring	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

(4) Femoral artery sheath removal	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) MAP	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(6) PA/Swan-Ganz	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(7) PCW pressure	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(8) PVR	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(9) Radial a-line	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(10) SVO2	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(11) SVR	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Intra aortic balloon pump	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Monitoring				
(1) 12 lead EKG interpretation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Arrhythmia interpretation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Lead placement	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) Rhythm strip assessment	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) Set up and run 12 lead EKG	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Pacemaker				
(1) External	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Permanent	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Temporary	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) Transthoracic (epicardial)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Ventricular assist device (RVAD or LVAD)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4. Care of the patient with:				
a. Abdominal aortic aneurysm repair	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Acute MI	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Cardiac arrest	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Cardiac tamponade	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Congestive heart failure (CHF)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. EP study & ablation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Heart transplant	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Immediate post open-heart surgery	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Infective endocarditis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Myocardial contusion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Pericarditis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Post AICD insertion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
m. Post arthroectomy (DCA)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
n. Post commissurotomy, valve repair, valve replacement	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
o. Post intracoronary stent placement	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
p. Post percutaneous balloon valvuloplasty	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
q. Post rotoblade	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
r. Pre/post angioplasty	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
s. Pre/post cardiac cath	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
5. Medications				
a. Amiodarone (Cardarone)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Atropine	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Bicarbonate	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Bretylium (Bretylol)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Digoxin (Lanoxin)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Diltiazem (Cardizem)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Dobutamine (Dobutrex)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Dopamine (Intropin)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Epinephrine (Adrenalin)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

j. Esmolol (Brevibloc)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Inocor (Amrinone)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Lidocaine (Xylocaine)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
m. Metoprolol (Lopressor)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
n. Nipride (Nitroprusside)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
o. Nitroglycerine (Tridil)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
p. Procainamide (Pronestyl)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
q. Reteplase recombinant (Retavase)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
r. Streptokinase	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
s. TPA (Alteplase)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
t. Verapamil (Calan, Isoptin, Verelan)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

B. PULMONARY

1. Assessment				
a. Adventitious breath sounds	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Rate and work of breathing	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2. Interpretation of lab results - arterial blood gases	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3. Equipment & procedures				
a. Air leak troubleshooting				
(1) Mediastinal chest tube removal	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Pleural chest tube removal	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Airway management devices/suctioning				
(1) Endotracheal tube/suctioning	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Extubation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Nasal airway/suctioning	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) Oximetry	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) Sputum specimen collection	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(6) Tracheostomy/suctioning	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Assist with				
(1) Bronchoscopy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Chest tube insertion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Emergency tracheostomy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) Thoracentesis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Establishing an airway				
(1) Assist with intubation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Oral airway insertion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Identification/intervention for respiratory complications				
(1) Aspiration	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) Laryngospasm	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Tension pneumothorax	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) Use of Pleurevac or Thoraclex drainage	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) Use of water seal drainage	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. O2 therapy & medication delivery systems				
(1) Ambu bag and mask	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) ET tube	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(3) Face masks	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) Nasal cannula	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) Portable O2 tank	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(6) Trach collar	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Ventilator management				
(1) External CPAP	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(2) High frequency jet ventilation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

(3) IMV	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(4) PEEP	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(5) Pressure support	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
(6) Weaning modes & T-piece weaning	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4. Care of the patient with:				
a. Acute pneumonia	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. ARDS	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Chest trauma	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. COPD	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Cor pulmonale	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Fresh tracheostomy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Lobectomy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Lung transplant	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Near drowning	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Pneumonectomy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Pulmonary edema/hypertension	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Pulmonary embolism	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
m. Status asthmaticus	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
n. Thoracotomy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
o. Tuberculosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
5. Medications				
a. Alupent (Metaproterenol)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Aminophylline (Theophylline)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Bronkosol (Isoetharine hydrochloride)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Corticosteroids	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Ventolin (Albuterol)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

C. NEUROLOGICAL

1. Assessment				
a. Cranial nerves	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Glasgow coma scale	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Level of consciousness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Pathologic reflexes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Reflex/motor deficits	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Visual or communication deficits	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2. Equipment & procedures				
a. Assist with lumbar puncture	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Halo traction/cervical tongs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Intracranial pressure monitoring	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Nerve stimulators	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Rotating bed	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Seizure precautions	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Spinal precautions	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Stryker frame	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Use of hyper/hypothermia blanket	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3. Care of the patient with:				
a. Aneurysm precautions	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Basal skull fracture	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Closed head injury	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Coma	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. CVA	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. DTs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

- g. Encephalitis
- h. Externalized VP shunts
- i. Increased ICP
- j. Laminectomy
- k. Meningitis
- l. Metastatic tumor/intracranial tumor resection
- m. Multiple sclerosis
- n. Post craniotomy
- o. Spinal cord injury
- p. Ventriculostomy
- 4. Medications
 - a. Barbiturate induced coma
 - b. Decadron (Dexamethasone)
 - c. Dilantin (Phenytoin)
 - d. Epidural administration
 - e. Phenobarbital
 - f. Valium (Diazepam)

D. GASTROINTESTINAL

- 1. Assessment
 - a. Abdominal/bowel sounds
 - b. Nutritional
- 2. Interpretation of lab results
 - a. Serum ammonia
 - b. Serum amylase
 - c. LFTs
- 3. Equipment & procedures
 - a. Administration of tube feeding
 - b. Balloon tamponade (Sengstaken Blakemore)
 - c. Feeding pump
 - d. Flexible feeding tube (i.e., Corpak, Dobhoff)
 - e. Gravity feeding
 - f. Iced saline lavage
 - g. Management of
 - (1) Gastrostomy tube
 - (2) Jejunostomy tube
 - (3) T-tube
 - (4) TPN and lipids administration
 - (5) PPN (peripheral parenteral nutrition)
 - h. Placement of nasogastric tube
 - i. Salem sump to suction
- 4. Care of the patient with:
 - a. Blunt trauma
 - b. Bowel obstruction
 - c. Colostomy
 - d. ERCP
 - e. Esophageal bleeding
 - f. GI bleeding
 - g. GI surgery
 - h. Hepatitis
 - i. Ileostomy
 - j. Inflammatory bowel disease

- k. Liver failure
- l. Liver transplant
- m. Pancreatitis
- n. Paralytic ileus
- o. Penetrating trauma
- 5. Medications
 - a. AquaMephyton (Vitamin K)
 - b. Inderal (Propranolol)
 - c. Kayexelate
 - d. Lactulose (Cephulac)
 - e. Pitressin (Vasopressin)

E. RENAL/GENITOURINARY

- 1. Assessment
 - a. A-V fistula/shunt
 - b. Fluid status
- 2. Interpretation of lab results
 - a. BUN & creatinine
 - b. Serum electrolytes
- 3. Equipment & procedures
 - a. Bladder irrigation
 - b. Insertion & care of straight and Foley catheter
 - (1) 3-way Foley
 - (2) Female
 - (3) Male
 - c. Supra-pubic
- 4. Care of the patient with:
 - a. Acute renal failure
 - b. CAVH dialysis
 - c. Hemodialysis
 - d. Nephrectomy
 - e. Peritoneal dialysis
 - f. Renal rejection syndrome
 - g. Renal transplant
 - h. TURP
 - i. Urinary diversion (ileal conduit nephrostomy)
 - j. Urinary tract infection

F. ENDOCRINE/METABOLIC

- 1. Interpretation of lab results
 - a. Blood glucose
 - b. Thyroid studies
- 2. Equipment & procedures
 - a. Blood glucose measuring device: Type : _____
 - b. Blood glucose monitoring
 - c. Performing finger stick
- 3. Care of the patient with:
 - a. Diabetes mellitus
 - b. Diabetic ketoacidosis
 - c. Disorders of adrenal gland (e.g., Addison's disease)
 - d. Disorders of pituitary gland (e.g., DI)
 - e. Drug overdose

- | | | | | |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| f. Hyperthyroidism (Grave's disease) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| g. Hypothyroidism | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| h. Insulin shock | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| i. Thyroidectomy - disorders of thyroid gland | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| 4. Medication - insulin pump | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

G. WOUND MANAGEMENT

- | | | | | |
|------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 1. Assessment | | | | |
| a. Skin for impending breakdown | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| b. Stasis ulcers | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| c. Surgical wound healing | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| 2. Equipment & procedures | | | | |
| a. Air fluidized, low airloss beds | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| b. Sterile dressing changes | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| c. Wound care/irrigations | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| 3. Care of the patient with: | | | | |
| a. Burns | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| b. Pressure sores | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| c. Staged decubitus ulcers | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| d. Surgical wounds with drain(s) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| e. Traumatic wounds | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

H. PHLEBOTOMY/IV THERAPY

- | | | | | |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| 1. Equipment & procedures | | | | |
| a. Administration of blood/blood products | | | | |
| (1) Cryoprecipitate | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (2) Packed red blood cells | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (3) Plasma/albumin | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (4) Whole blood | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| b. Drawing blood from central line | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| c. Drawing venous blood | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| d. Starting IVs | | | | |
| (1) Angiocath | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (2) Butterfly | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (3) Heparin lock | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| 2. Care of the patient with: | | | | |
| a. Central line/catheter/dressing - | | | | |
| (1) Broviac | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (2) Groshong | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (3) Hickman | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (4) Portacath | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| (5) Quinton | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| b. Peripheral line/dressing | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

I. PAIN MANAGEMENT

- | | | | | |
|---------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 1. Assessment of pain level/tolerance | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| 2. Care of the patient with: | | | | |
| a. Epidural anesthesia/analgesia | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| b. IV conscious sedation | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| c. Patient controlled analgesia | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

J. MISCELLANEOUS

- | | | | | |
|------------------------------|--|--|--|--|
| 1. Care of the patient with: | | | | |
|------------------------------|--|--|--|--|

- a. Anaphylactic shock
- b. Disseminated intravascular coagulation (DIC)
- c. Hypovolemic shock
- d. Multi-system organ failure
- e. Organ/tissue donation
- f. Septic shock

AGE SPECIFIC PRACTICE

A. Newborn/Neonate (birth - 30 days)	D. Preschooler (3 - 5 years)	G. Young adults (18 - 39 years)
B. Infant (30 days - 1 year)	E. School age children (5 - 12 years)	H. Middle adults (39 - 64 years)
C. Toddler (1 - 3 years)	F. Adolescents (12 - 18 years)	I. Older adults (64+)

EXPERIENCE WITH AGE GROUPS

- | | A | B | C | D | E | G | H | I |
|--|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Able to adapt care to incorporate normal growth and development. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Able to adapt method and terminology of patient instructions to their age, comprehension and maturity level. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Can ensure a safe environment reflecting specific needs of various age groups. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

My experience is primarily in: (Please indicate number of years)

- | | | |
|--|--|---|
| <input type="radio"/> Medical _____ year(s) | <input type="radio"/> Cardiothoracic _____ year(s) | <input type="radio"/> Neuro _____ year(s) |
| <input type="radio"/> Neurological _____ year(s) | <input type="radio"/> Cardiovascular _____ year(s) | <input type="radio"/> Burn _____ year(s) |
| <input type="radio"/> Trauma _____ year(s) | <input type="radio"/> Coronary care _____ year(s) | <input type="radio"/> PACC _____ year(s) |
| <input type="radio"/> Other (specify) _____ | _____ year(s) | |

The information I have given is true and accurate to the best of my knowledge. I hereby authorize Professional Nursing Service to release Critical Care/Intensive Care Skills Checklist to client facilities of PNS in relations to consideration of employment as a Traveler with those facilities.

Signature

Date

Address

Phone